

First aid procedures

Review of what is different between the typical CSP protocols and the COVID-19 modified protocols.

- Basic Life Support (AR and CPR)

Basic Life Support (AR and CPR)

DRAFT

Extra measures must be taken to provide Basic Life Support in a COVID-19 world. One of the most stunning modification to our protocols is the removal of ventilations from the entire Artificial Respiration and Cardiopulmonary Resuscitation.

Based on the latest directives and discussions between CSP Yukon Zone instructors on October 16th (*BLS non covid vs covid procedures summary.doc*):

		Non COVID (normal)	
time	mod	Check No change	
am	BLS	vital etc	

		Position of airway, jaw Clearing the airway as obstruction, etc, etc...
		Proper seal, correct equipment Re- (mask, breather A.R. rebreather, etc) with triangular , only rate etc, etc ...
		How, type Same no but blind must use wear Suction leaving, that surgical 5 mask, seconds, shield/eye etc pro etc..

		<p>Fit, size Same when how must inf wear OPAs gown, child rigid positioning, etc, faceshield/eye etc pro ...</p>
		<p>How to set up, O2 rate, fit for DO BVM, CNOT 1 USE or 2 person, points of failure (physical/mechanical)</p>

PM

Call
ems
as of
needed,
engage
with
patient,
ab NOT
thru
(back
in close
infant),
size
of to
pers
pres
wor
chest
airway
Choking
A, C
go
unconscious
(look,
listen
and
feel),
attempt
vent,
chest
compressions
(back
and
chest
in
infant)
at
30:2,
etc,
etc,...

All
above
except...
Do,
NOT
use
face
theirs
sister
look
and
feel
only

After
30
chest
compressions
look
and
feel,
attempt
but
DO
NOT
VENTILATE
use
nasal
prongs
(cannula)
only

		Follow instructions, proper padNO placechange, AEDexcept use NO in VENTILATIONS certain locations, rebreather CPR with O2 between AED only checks, etc, etc,...
		NO change Concussion rate NO CPR(30:2), A,C,position, etc rebreather etc. with O2 only
	CPR test	

...

	Traditional world	COVID-19 world
Artificial Respiration (AR)		
Suction		
Use of Oropharyngeal airway (OPA)		

Bag Valve Mask		